

developed dark fuscous bars. Hindwings grey-whitish, thinly scaled, suffused grey posteriorly; cilia grey-whitish.

BRAZIL, Parintins, October (*Parish*); 3 ex.

CRASIMORPHA, n. g.

Head with appressed scales; ocelli posterior; tongue developed. Antennæ $\frac{4}{5}$, basal joint moderate, without pecten. Labial palpi rather long, curved, ascending, second joint thickened with dense scales, with rough projecting scales towards apex beneath, terminal joint shorter than second, rather stout, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiæ with rough hairs above. Forewings with some raised scales; 1 *b* furcate, 2 from towards angle, 3 and 4 stalked, 7 and 8 stalked, 7 to costa, 6 and 9 approximated to 8 at base, 11 from middle. Hindwings over 1, trapezoidal-ovate, termen not sinuate, cilia $\frac{1}{3}$; 3 and 4 short-stalked, 5 rather approximated, 6 and 7 stalked.

Close to *Oestomorpha*.

Crasimorpha peragrata, n. sp.

♀. 23 mm. Head white sprinkled grey on crown. Palpi white, two bands of second joint and base of terminal dark fuscous. Thorax white sprinkled grey, patagia tinged grey. Forewings elongate, apex rounded, termen rounded, somewhat oblique; light fuscous, mostly irregularly overlaid white suffusion, especially towards costal and dorsal areas, a few scattered blackish scales; a slender irregular blackish streak along fold mostly edged white beneath, and a similar longitudinal streak from middle of disc to termen, a similar streak above posterior portion of this; a blackish longitudinal streak between these two towards termen, and sub-dorsal one on posterior half; an elongate dark fuscous mark on middle of costa; cilia white barred light fuscous. Hindwings and cilia grey.

FRENCH GUIANA, St. Jean, R. Maroni, July; 1 ex.

Sophronia sagittans, n. sp.

♂ ♀. 13-14 mm. Head whitish-ochreous, face more whitish. Palpi white, second joint light greyish-ochreous except apical edge, tuft long, rough. Thorax light ochreous-brown, sometimes dorsally whitish-tinged. Forewings rather narrow, pointed, termen slightly sinuate, very oblique; ochreous-fulvous or ochreous-brown; a white streak, attenuated basally, along costa from base to near middle, thence leaving costa and narrowed to beyond middle; a very oblique white striga from costa at $\frac{2}{3}$, near termen acutely angulated to tornus, edged posteriorly with dark grey speckling strongest in disc, and preceded in angle by a fine black dash; two or sometimes three inwards-oblique white marks on costa posteriorly, followed by

two black marks before apex : cilia whitish with lines of dark grey specks, above apex with stronger dark grey speckling on basal half and grey outwardly, white on costal marks. Hindwings grey; cilia light greyish.

PALESTINE, Haifa, May (*Buxton*); 3 ex. (type Brit. Mus.). Nearest *acaudella* and *finitimella*, which, however, both have a white dorsal streak.

Sophronia catharurga, n. sp.

♀. 11 mm. Head silvery-white. Palpi wholly white, tuft moderate. Thorax grey, anteriorly suffused white. Forewings rather narrow, pointed, termen very oblique; greyish-ochreous, on costa posteriorly sprinkled dark grey, dorsal area broadly suffused silvery-whitish; a moderate shining white costal streak from base to middle, thence leaving costa and continued narrowed to $\frac{4}{5}$, where it receives an oblique white strigula from costa at $\frac{2}{5}$; beneath this three short longitudinal white lines in disc from middle, uppermost running into costal streak, some white irroration beyond these; a spot of whitish suffusion on tornus, followed by some dark grey irroration: cilia whitish, a dark fuscous subbasal line, on costa also a basal line, tips fuscous, a dark fuscous apical spot. Hindwings grey-whitish; cilia whitish.

PALESTINE, Akka, June (*Buxton*); 1 ex. (Brit. Mus.).

CROSSOBELA, n. g.

Head with appressed scales; ocelli posterior; tongue developed. Antennæ $\frac{4}{5}$, in ♂ serrulate, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint thickened with dense rough projecting scales beneath throughout, above with scales rather expanded towards apex, terminal joint half second, slender, pointed. Maxillary palpi rudimentary. Forewings 2-6 separate, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, termen emarginate, cilia 2; 3 and 4 rather approximated, 5 remote, 6 and 7 stalked.

Allied to *Brachyacma*, differing by second joint of palpi rough-scaled beneath.

Crossobela barysphena, n. sp.

♂. 11-12 mm. Forewings elongate, apex pointed, termen very obliquely rounded; light yellow-ochreous, somewhat sprinkled light grey; a dark fuscous thinly whitish-sprinkled costal streak from $\frac{1}{4}$ to apex, attenuated anteriorly and becoming broad posteriorly; a narrow dorsal similar streak, extended as a slender terminal streak to apex; discal stigmata moderate, blackish: cilia light grey, on costa dark grey. Hindwings rather dark grey; cilia light grey.

CYPRUS, Limassol, February (*Mavromoustaki*); 2 ex. (Brit. Mus.), unfortunately much damaged in transit, but determinable.

OXYSACTIS, n. g.

Essential characters of *Brachyacma*, but hindwings with cubital pecten.

Type *sciritis* Meyr.

RHYNCHOTONA, n. g.

Head with appressed scales; ocelli posterior; tongue absent. Antennæ $\frac{3}{4}$, in ♂ serrulate, shortly ciliated, basal joint elongate, without pecten. Labial palpi extremely long, porrected, second joint thickened with dense scales loosely appressed and projecting at apex, terminal joint short, slender, acute, projecting little from apical scales of second. Maxillary palpi obsolete. Posterior tibiæ shortly rough-scaled above. Forewings 2 and 3 stalked from angle, 7 to just above apex, 8 and 9 out of 7, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex rather produced, termen sinuate, cilia 1; with cubital pecten; 3-5 nearly approximated at base, 6 and 7 connate.

Perhaps a derivative of *Dichomeris*.

Rhynchotona phaeostrota, n. sp.

♂. 10 mm. Head, palpi, thorax greyish-ochreous. Forewings elongate, slightly dilated, apex obtuse-pointed, termen obliquely rounded; greyish-ochreous speckled dark fuscous; an indistinct small cloudy dark fuscous spot in disc at $\frac{1}{4}$; plical and second discal stigmata small, blackish: cilia greyish-ochreous (imperfect). Hindwings rather dark grey; cilia grey.

PERU, Jurimaguas, March (*Parish*); 1 ex.

Dichomeris neatodes, n. sp.

♂. 20 mm. Head, thorax whitish-grey. Palpi white, second joint dark fuscous except apex and apical hairs, with moderate tuft, terminal joint anteriorly blackish-sprinkled. Antennæ serrulate, ciliations 1. Forewings elongate, termen rounded, rather strongly oblique; light grey, suffusedly mixed whitish towards costa, with indistinct fine lines of blackish scales on veins, and indistinct fine lines of whitish scales between these posteriorly and towards dorsum, within cell with obscure streaks of blackish irroration above lower edge anteriorly and beneath upper edge posteriorly; stigmata small, black, first discal indistinct, plical linear, rather before first discal; a marginal series of rather large triangular black dots on apical part of costa and termen: cilia grey-whitish with three grey lines. Hindwings light grey; cilia grey-whitish.

CYPRUS, Platres, 5000-6500 feet, September (*Hayward*); 1 ex. Perhaps allied to *juniperella*.

Sarisophora ptochomorpha, n. sp.

♀. 14 mm. Head violet-fuscous, face and sides of crown ochreous-yellowish. Palpi whitish-ochreous, second joint dark fuscous except apex, anterior edge of terminal joint dark fuscous. Antennæ pale yellowish ringed dark fuscous. Thorax dark violet-fuscous. Forewings termen slightly rounded, rather oblique; 7 to apex, 9 separate; grey suffusedly mixed dark violet-fuscous; discal stigmata obscurely darker: cilia grey, basal half mixed dark fuscous. Hindwings rather dark grey; cilia grey.

CYPRUS, Platres, July (*Hayward*); 1 ex. The European *briantilla*, *pallicornella*, and *flavissimella* are all referable to *Sarisophora* and not to *Lecithocera*.

NEOCORODES, n. g.

Head loosely scaled, sidetufts raised, connivent; ocelli posterior; tongue developed. Antennæ 1, in ♂ rather stout, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint slightly longer, slender, acute. Maxillary palpi rudimentary. Forewings 2 from towards angle, 3 and 4 approximated from angle, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal-ovate, cilia over 1; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Near *Achoria* and *Enthetica*.

Neocorodes amnesta, n. sp.

♂. 10 mm. Head ochreous-whitish, central stripe greyish. Palpi whitish, second joint greyish except apex, anterior edge of terminal joint grey. Antennæ ochreous-whitish ringed fuscous. Forewings elongate, apex pointed, termen very obliquely rounded; light fuscous irrorated dark fuscous; discal stigmata forming small cloudy dark fuscous spots: cilia whitish-grey, base sprinkled dark fuscous. Hindwings grey; cilia whitish-grey.

CYPRUS, Platres, 5000-6000 feet, September (*Hayward*); 1 ex.

Mnesistega convexa, n. sp.

♀. 18-20 mm. Head pale yellow-ochreous, base of sidetufts dark fuscous. Palpi whitish-ochreous, second joint dark fuscous except towards apex. Thorax dark purplish-fuscous. Forewings elongate, costa gently arched, apex obtuse-pointed, termen very obliquely rounded; light fulvous-ochreous, slightly sprinkled dark fuscous, costa more irrorated dark fuscous, more strongly and suffusedly towards base; a moderate dark purple-fuscous basal patch, widened upwards but not reaching costa; plical and first discal stigmata cloudy, rather dark fuscous, plical rather obliquely posterior; second discal and a spot on dorsum united to form an

irregular nearly direct rather dark purplish-fuscous streak: cilia pale ochreous, some fuscous specks near base. Hindwings light ochreous-grey; cilia concolorous, base pale.

ASSAM, Shillong, Cherrapunji, September (*Fletcher*); 4 ex. Very like *talantodes*, but forewings much broader and costa more arched.

SYNCATHEDRA, n. g.

Head with appressed scales, sidetufts rather loose; ocelli posterior; tongue developed. Antennæ $\frac{5}{6}$, somewhat flattened, not tapering, in ♂ stout, simple, in ♀ rather slender, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, with appressed scales, terminal joint shorter than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiæ clothed with rough hairs above. Forewings 1 *b* furcate, 2 and 3 stalked from angle, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, elongate-ovate, tolerably pointed, cilia 1; 3 and 4 connate, 5 nearly parallel, 6 and 7 long-stalked.

Syncathedra criminata, n. sp.

♂ 15–20 mm., ♀ 18–25 mm. Head and thorax ♂ grey mixed dark fuscous or wholly dark fuscous, ♀ fuscous mixed whitish. Palpi whitish suffusedly irrorated dark fuscous except base and apex of terminal joint. Antennæ ♂ dark grey or dark fuscous, ♀ fuscous speckled whitish, a dark fuscous apical band, last apical joint whitish. Forewings elongate, apex pointed, termen ♂ nearly straight, ♀ sinuate, very oblique; ♂ dark fuscous, ♀ fuscous with bases of scales more or less whitish; more or less developed streaks of coarse black irroration between veins; stigmata forming suffused irregular black spots, plical rather beyond first discal: cilia ♂ dark grey, ♀ grey towards base mixed whitish and fuscous. Hindwings ♂ dark fuscous, cilia dark grey; ♀ rather dark grey, cilia grey, base whitish.

ASSAM, Cherrapunji (*Fletcher*); 61 ex. (33 ♂, 28 ♀).

STEREMNIODES, n. g.

Head with appressed scales; ocelli posterior; tongue developed. Antennæ $\frac{4}{5}$, in ♂ stout, serrulate, simple, basal joint moderate, without pecten. Labial palpi moderately long, curved, ascending, second joint much thickened with dense appressed scales, terminal joint shorter than second, stout, pointed. Maxillary palpi rudimentary. Posterior tibiæ rough-haired above. Forewings 2 and 3 stalked from angle, 4 approximated, 5 absent, 6 from near 7, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen faintly sinuate, cilia $\frac{3}{5}$; 3 and 4 stalked, 5 rather approximated, 6 and 7 stalked.

Steremniodes sciactis, n. sp.

♂. 16 mm. Head, palpi, thorax fuscous. Abdomen grey, segment 2 whitish-ochreous dorsally, anal tuft whitish-grey. Forewings elongate, termen nearly straight, somewhat oblique; fuscous, irregularly irrorated dark fuscous tending to form suffused dark fuscous streaks between veins and along fold, costal area suffused dark fuscous; small dark fuscous marginal dots round apex and termen: cilia grey, base and tips pale greyish-ochreous. Hindwings grey speckled darker; cilia grey, base pale greyish-ochreous.

BRAZIL, Teffé, January (*Parish*); FRENCH GUIANA, La Forestière, June; 2 ex.

Enthetica tribrachia, n. sp.

♂ ♀. 14–15 mm. Head whitish-ochreous. Palpi whitish, second joint fuscous except apex, anterior edge of terminal joint dark fuscous. Antennæ whitish-ochreous. Thorax whitish-ochreous, shoulders infuscated. Forewings narrow, apex pointed, termen extremely obliquely rounded; ♂ light brownish-ochreous, ♀ fuscous; stigmata forming elongate dark fuscous marks, plical obliquely before first discal: cilia pale ochreous or greyish-ochreous. Hindwings 3 and 4 connate; pale grey; cilia as in forewings.

S. INDIA, Palnis, 7000 feet, August (*Campbell, Fletcher*); 2 ex.

Lecithocera opportuna, n. sp.

♂ ♀. 16–17 mm. Head whitish-ochreous. Palpi whitish, second joint fuscous except apex, anterior edge of terminal joint dark fuscous. Antennæ whitish-ochreous or whitish, dotted or ringed fuscous. Thorax light fuscous. Posterior tibiæ dark grey, tip whitish. Forewings apex obtuse-pointed, termen concave, oblique; 8 and 9 out of 7; light fuscous, sometimes darker posteriorly; discal stigmata dark fuscous, an additional dot beneath second; an oblique ochreous-whitish mark from costa at $\frac{3}{4}$, whence a faint pale line curved or bent above middle runs to dorsum at $\frac{5}{8}$: cilia dark grey, tips ochreous-whitish. Hindwings 4 absent, 3 and 5 connate; pale greyish, more or less tinged whitish-ochreous; cilia whitish-ochreous, two more or less faint greyish shades.

ASSAM, Cherrapunji (*Fletcher*); 6 ex.

Lecithocera xanthoteles, n. sp.

♀. 18–20 mm. Head ochreous-yellow. Palpi ochreous-yellow, anterior edge of terminal joint dark fuscous. Antennæ pale yellowish above, fuscous beneath. Thorax dark purple-fuscous. Abdomen dark fuscous, apical third ochreous-yellow. Forewings slightly dilated, apex obtuse, termen hardly sinuate, rather oblique; 3 and 4 out of 2, 8 and 9 out of 7; dark purple-fuscous: cilia

dark fuscous. Hindwings dark fuscous; cilia fuscous, a darker subbasal shade.

ASSAM, Sadiya, May (*Fletcher*); BURMA, Rangoon, August; 2 ex.

Lecithocera dirupta, n. sp.

♂. 15 mm. Head ochreous-white. Palpi whitish, second joint fuscous except apex, anterior edge of terminal joint dark fuscous. Antennæ ochreous-whitish. Thorax ochreous-whitish, edge of shoulder dark fuscous. Forewings rather narrow, slightly dilated, apex obtuse-pointed, termen nearly straight, very oblique; 7 and 8 stalked, 7 to termen; ochreous-whitish, slightly sprinkled fuscous; base of costa dark fuscous; a minute black strigula beneath costa near base; stigmata blackish, discal forming moderate spots, plical forming a streak extending from beneath first discal halfway to base, an elongate spot beneath second discal, and extending anteriorly before it; some irregular fuscous suffusion round these markings, and veins between cell and termen more or less streaked with dark fuscous suffusion, especially 7 and 8; an irregular terminal line of dark fuscous suffusion: cilia ochreous-whitish, two faint fuscous lines, at apex a spot of dark fuscous suffusion. Hindwings whitish-grey; cilia ochreous-whitish, two faint fuscous shades.

S. INDIA, Palnis, Kodaikanal, 7000 feet, August (*Fletcher*); 1 ex.

Lecithocera carcerata, n. sp.

♂. 15 mm. Head, antennæ whitish-ochreous. Palpi whitish, second joint fuscous except towards apex, anterior edge of terminal joint fuscous. Thorax light lilac-fuscous. Forewings slightly dilated, apex pointed, termen sinuate, oblique; 8 and 9 out of 7, 7 to termen; light lilac-fuscous; a dark fuscous transverse fasciate blotch from dorsum at $\frac{2}{5}$ reaching $\frac{3}{4}$ across wing, both sides prominent in middle; an elongate blackish mark on costa beyond this; two dark fuscous dots transversely placed in disc at $\frac{3}{5}$; a nearly straight ochreous-whitish subterminal line at $\frac{4}{5}$, preceded by a fascia of rather dark fuscous suffusion; some minute blackish terminal dots: cilia whitish-ochreous, tinged grey on termen. Hindwings light grey; cilia pale greyish.

ASSAM, Shillong, July (*Fletcher*); 1 ex.

Lecithocera recurvata, n. sp.

♀. 16-18 mm. Head pale fuscous, face whitish. (Palpi broken.) Antennæ whitish, slenderly ringed fuscous towards base. Thorax light fuscous. Posterior tibiæ dark grey, apex white. Forewings apex pointed, termen concave, rather oblique; 7 absent, 8 and 9 stalked; light fuscous; terminal edge dark fuscous: cilia white, on costa and beneath tornus light grey. Hindwings dark grey; cilia greyish.

PHILIPPINES, Mindanao, Baco R., February (*Mounsey*); 2 ex.

Lecithocera ochrocapna, n. sp.

♀. 21 mm. Head, thorax whitish-grey-ochreous. Palpi whitish, second joint grey except tip. Antennæ whitish, basal $\frac{2}{5}$ ringed fuscous. Posterior tibiæ grey, apical fifth white. Forewings somewhat dilated, apex obtuse-pointed, termen hardly sinuate, oblique; 2 and 4 stalked, 3 absent, 7 absent, 8 and 9 stalked; ochreous-fuscous; a small suffused ochreous-whitish oblique wedgeshaped mark on costa at $\frac{4}{5}$; cilia ochreous-brownish, on termen ochreous-whitish with a dark fuscous basal line becoming thicker round tornus. Hindwings grey; cilia grey, a darker sub-basal shade.

PHILIPPINES, Mindanao, Mt. Apo, 6500 feet, May (*Mounsey*). Near *orbata*, which has posterior tibiæ wholly grey.

Lecithocera alcestis, n. sp.

♂ ♀. 13-16 mm. Head, thorax pale greyish-ochreous. Palpi whitish, second joint grey except tip, anterior edge of terminal joint dark grey. Antennæ whitish ringed fuscous. Posterior tibiæ dark grey, apical half white. Forewings hardly dilated, apex obtuse-pointed, termen sinuate, rather oblique; 8 and 9 out of 7; light fuscous; a cloudy ochreous-whitish dot on costa at $\frac{4}{5}$; cilia fuscous, on termen outer half beyond a blackish-grey antemedian shade ochreous-whitish. Hindwings grey, paler anteriorly; cilia pale greyish.

COORG, Dibidi, 3500 feet, January (*Newcome*); KANARA, Anshi, December (*Maxwell*); 2 ex.

Lecithocera affusa, n. sp.

♀. 16 mm. Head light violet-grey, sides whitish-ochreous. Palpi whitish-ochreous, second joint tinged yellow, anterior edge of terminal joint dark grey. Antennæ whitish-ochreous ringed grey. Thorax light violet-grey. Forewings slightly dilated, apex obtuse, termen obliquely rounded; 4 and 5 stalked, 8 and 9 out of 7; pale ochreous slightly speckled grey; discal stigmata small, blackish-grey; cilia whitish-ochreous. Hindwings rather dark grey; cilia whitish-grey.

ASSAM, Khasis, June; 1 ex. Allied to *choritis*.

Lecithocera cassiterota, n. sp.

♀. 16 mm. Head shining violet-grey, orange lateral lines. Palpi orange, terminal joint dark fuscous anteriorly. Antennæ yellowish. Thorax violet-grey. Forewings hardly dilated, apex obtuse-pointed, termen concave, rather oblique; 3 absent, 4 and 5 connate, 7 absent, 8 and 9 stalked; glossy violet-grey; a short pale metallic-blue streak on base of costa; markings orange, irregularly edged blackish; irregular subcostal and median streaks from base terminating in a direct transverse antemedian fascia; a rather

oblique postmedian fascia, whence an irregular costal streak runs to near apex, and a median streak to termen; a fascia of blackish suffusion mixed orange crossing from apex of costal streak to lower part of postmedian fascia; a blackish marginal line round apex and termen except on median streak: cilia ochreous-yellow, outer half pale shining grey. Hindwings grey; cilia light grey.

PHILIPPINES, Manila; 1 ex.

Lecithocera hippotarcha, n. sp.

♀. 19 mm. Head dark violet-fuscous, orange lateral lines. Palpi second joint thickened and somewhat roughened beneath towards apex, yellow-ochreous with broad dark fuscous posterior streak and mixed dark fuscous on anterior edge, terminal joint whitish with black lateral lines. Antennæ dark fuscous, thickened with rather rough scales on basal half, apical fourth pale ochreous. Thorax dark purple-fuscous. Forewings not dilated, apex obtuse-pointed, termen sinuate, rather oblique; 3 absent, 8 and 9 out of 7; dark purple-fuscous; some undefined greyish-ochreous suffusion towards costa before middle; an elongate wedge-shaped yellow-ochreous spot on costa at $\frac{2}{3}$; a small yellow-ochreous dot on tornus: cilia dark purplish-fuscous. Hindwings 3 absent; dark bronzy-grey; cilia dark grey.

CEYLON, Kandy, September (*Mackwood*); 1 ex.

Lecithocera melitopyga, n. sp.

♀. 18 mm. Head, palpi ochreous-yellow (terminal joint broken). Antennæ yellow-whitish at base gradually becoming light grey, basal joint ochreous-yellow. Thorax dark purple-grey. Abdomen segment 1 dark grey, 2-5 dark brown above with yellow marginal rings and dark grey on sides, rest ochreous-yellow with base of segments brown dorsally. Forewings somewhat dilated, apex obtuse, termen faintly sinuate, rather oblique; 3 and 4 out of 2, 5 connate, 8 and 9 out of 7; dark purple-grey: cilia concolorous. Hindwings dark grey; cilia grey, lighter towards base.

CEYLON, May (*Alston*); 1 ex.

Lecithocera residua, n. sp.

♂. 23 mm. Head whitish, sides of face and a spot in middle of collar grey. (Palpi missing.) Thorax whitish, shoulders and a broad dorsal stripe suffused purplish-grey. Abdomen ochreous-whitish. Forewings slightly dilated, apex tolerably pointed, termen concave, little oblique; 2 and 4 stalked, 3 absent, 8 and 9 out of 7; ochreous-grey-whitish, slightly grey-sprinkled; an irregular-semioval purplish-fuscous blotch extending along dorsum from near base to $\frac{3}{5}$, edge prominent at $\frac{2}{5}$, where it reaches $\frac{3}{4}$ across wing; dorsal $\frac{3}{5}$ from this to termen mixed light grey suffusion and irregularly sprinkled darker fuscous, with two or three small spots

in disc beyond middle; a dark fuscous marginal line round termen and tornus and extending a little way up fold: cilia whitish, on termen outer half suffused light grey. Hindwings pale whitish-grey; cilia ochreous-whitish, outer half faintly tinged grey.

ASSAM, Khasis; 1 ex.

PLACANTHES, n. g.

Head smooth; ocelli very small, posterior; tongue developed. Antennæ probably over 1. Labial palpi long, recurved, second joint somewhat thickened, smooth, terminal joint longer than second, slender, acute. Maxillary palpi very short, loosely scaled. Posterior tibiæ shortly rough-scaled above. Forewings 1 *b* furcate, 2 and 4 stalked from angle, 3 absent, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, costa with some rough projecting scales anteriorly, apex forming a long pointed projection (nearly $\frac{1}{3}$ of wing), termen rectangularly emarginate beneath this, cilia $\frac{1}{5}$; 3 and 4 connate, 5 absent, 6 and 7 stalked.

Perhaps intermediate between *Philoptila* and *Crocantbes*.

Placanthes xanthomorpha, n. sp.

♂. 24 mm. Head, palpi ochreous-yellow (injured). Thorax dark fuscous, two posterior yellow spots. Forewings posteriorly dilated, apex obtuse-pointed, termen sinuate, rather oblique; blackish-fuscous, markings orange-yellow; an elongate-oblong spot above dorsum from near base to $\frac{1}{4}$; a small elongate spot beneath costa at $\frac{1}{5}$; a transverse blotch in disc before middle not reaching margins; a large oval blotch extending in disc from beyond middle to $\frac{5}{6}$, with small pointed posterior projection: cilia dark fuscous. Hindwings dark bronzy-fuscous; markings orange-yellow; a rather thick subcostal streak from $\frac{1}{4}$ to beyond middle; an irregular trapezoidal blotch occupying posterior half of dorsum and extending to near middle of disc; an oval blotch extending in disc from middle to $\frac{5}{6}$; a patch occupying apical projection except tip, including a row of scattered blackish scales, and extended as a pointed streak along upper part of termen; cilia dark grey.

PHILIPPINES, Mindoro, Laguna de Naujan, March (*Mounsey*); 1 ex.

Onebala coercita, n. sp.

♀. 12–13 mm. Head light violet-fuscous, sides tinged whitish. Palpi whitish, second joint fuscous except tip. Antennæ ochreous-whitish. Thorax dark fuscous, patagia whitish except shoulders. Forewings apex obtuse-pointed, termen somewhat sinuate, rather oblique; 2 and 3 very long-stalked, 4 and 5 short-stalked, 8 and 9 out of 7; violet-fuscous, slightly whitish-sprinkled; a broad streak of blackish-fuscous suffusion on basal fourth of costa, posteriorly suffused and undefined; an irregular-edged triangular blackish-

fuscous white-edged patch extending on dorsum from base to middle, apex reaching $\frac{2}{3}$ across wing near before middle, space between this and costal streak suffused whitish; a very oblique blackish white-edged strigula on costa before middle; second discal stigma forming a small blackish white-edged spot, sometimes a smaller similar spot beneath it; a suffused dark fuscous subterminal fascia of which discal portion is expanded into a broad blotch reaching second discal stigma, edged beneath and posteriorly by white scales, a small white mark on costa preceding this; a blackish terminal line: cilia whitish with two thick grey shades, on costa dark fuscous. Hindwings 4 and 5 short-stalked; light grey, thinly scaled and whitish-tinged anteriorly; cilia pale grey.

COORG, Dibidi, 3500 feet, February, April (*Newcome*); 2 ex.

Onebala foederalis, n. sp.

♂ ♀. 20-21 mm. Head, thorax fuscous sprinkled whitish. Palpi white, second joint irrorated dark fuscous, terminal joint slightly sprinkled dark fuscous. Forewings apex obtuse-pointed, termen faintly sinuate, oblique; 4 and 5 coincident, 8 and 9 out of 7; brownish, irregularly irrorated dark fuscous; stigmata forming small irregular dark fuscous spots, sometimes partially edged whitish, plical obliquely beyond first discal, these sometimes united and confluent with dark fuscous suffusion extending to costa at $\frac{1}{3}$; a whitish spot on costa at $\frac{3}{4}$, whence a whitish line, usually nearly obsolete but sometimes distinct, sharply angulated first inwards and then outwards, runs to tornus; two minute whitish dots on costa towards apex; sometimes scattered white scales on apical area: cilia light fuscous, base whitish, a dark fuscous subbasal shade. Hindwings 4 and 5 stalked; light grey, anteriorly whitish-tinged; cilia ochreous-whitish, a faint greyish subbasal shade.

S. INDIA, Nilgiris, Pykara, 7000 feet, March (*Andrewes*); 4 ex.

Onebala pogonias, n. sp.

♂. 18 mm. Head, thorax fuscous sprinkled ochreous-whitish. Palpi ochreous-whitish, irregularly irrorated dark fuscous, second joint with broad tuft of long rough hairscales beneath. Forewings apex obtuse-pointed, termen sinuate, very oblique; 4 and 5 coincident, 8 and 9 out of 7; purplish-fuscous irregularly irrorated darker, scaletips minutely whitish; stigmata forming small irregular dark fuscous spots, plical rather beyond first discal, connected with it and with costa at $\frac{1}{5}$ by oblique dark fuscous suffusion, edged posteriorly by an ochreous-whitish streak obtusely angulated on fold and reaching dorsum, second discal whitish-ringed; a small ochreous-whitish spot on costa at $\frac{4}{5}$, whence an obscurely indicated line angulated first inwards and then outwards runs to tornus: cilia fuscous mixed darker and pale-speckled, base ochreous-whitish, on costa with two small dark spots. Hindwings 4 and 5 stalked;

grey; cilia ochreous-whitish, a well-marked dark grey subbasal shade.

S. INDIA, Nilgiris, 3500 feet, March (*Andrewes*); 1 ex. At first sight extremely like *fœderalis*, but the palpal tuft is unique in the genus; the more oblique and sinuate termen of forewings, and dark shade of cilia of hindwings are also good characters.

Onebala cataclina, n. sp.

♂. 21 mm. Head whitish-ochreous. Palpi whitish-ochreous, second joint dark fuscous except towards apex, terminal joint with fine dark fuscous lateral lines. Thorax whitish-fuscous, shoulders dark fuscous. Forewings apex obtuse-pointed, termen sinuate, rather oblique; 4 and 5 coincident, 8 and 9 out of 7; whitish-ochreous suffused light grey and sprinkled dark fuscous; a small blackish spot on base of costa; an irregular purple-blackish streak from $\frac{1}{4}$ of costa to middle of dorsum, thickest on costa, first discal and plical stigmata forming spots on this, second discal stigma forming a small blackish spot ringed whitish-ochreous; a pale ochreous spot on costa at $\frac{3}{4}$, whence an obscure shade obtusely angulated in middle runs to tornus, terminal area beyond this forming a light fuscous fascia speckled blackish; two pale ochreous dots on costa towards apex: cilia fuscous, base pale ochreous with short projecting teeth. Hindwings 4 and 5 connate; light grey; cilia ochreous-whitish, a faint grey subbasal shade.

CEYLON, Maskeliya (*Pole*); 1 ex.

Onebala leptobrocha, n. sp.

♂. 14 mm. Head, thorax glossy grey, a fine white line above eyes. Palpi second joint grey, tip white, a blackish subapical ring, terminal joint dark grey, base white. Forewings apex obtuse-pointed, termen slightly rounded, very oblique; 2 and 3 coincident, 4 and 5 stalked, 8 and 9 out of 7; brownish-fuscous, basal third suffused grey; markings bluish-leadен; an angulated transverse streak before middle, costa from this dark fuscous, cut by small white marks at middle and $\frac{4}{5}$, from first a line running to dorsum before tornus with a posterior loop line leaving it near costa and rejoining it near dorsum, from second a direct transverse line to tornus with slight outward sinus in middle; a black streak along apical part of costa and line along termen, within these a bluish-leadен streak: cilia light grey, a fuscous subbasal shade. Hindwings and cilia grey; 4 and 5 stalked.

COORÆ, Dibidi, 3500 feet, August (*Newcome*); 1 ex.

Onebala commodata, n. sp.

♀. 15 mm. Head, thorax light fuscous, face ochreous-whitish. Palpi fuscous, second joint with whitish subapical ring, terminal joint posteriorly whitish. Forewings apex obtuse-pointed, termen

slightly sinuate, rather oblique; 2 and 3 coincident, 4 and 5 stalked, 8 and 9 out of 7; rather dark bronzy-fuscous; an obscure cloudy whitish transverse line at $\frac{2}{5}$, groundcolour rather darker before it; a slight cloudy whitish transverse mark on costa at $\frac{4}{5}$, and a præ-tornal dot opposite; sometimes some faint whitish speckling towards termen; a black marginal line round apical part of costa and termen, thicker on costa: cilia whitish, with thick subbasal rather dark fuscous shade, base ochreous tinged, on costa and beneath tornus grey. Hindwings 4 and 5 stalked; grey; cilia grey, base whitish.

CEYLON, Maturatta, March (*Pole*); 2 ex.

Onebala leucograptæ, n. sp.

♀. 15 mm. (Head and palpi injured.) Forewings apex tolerably pointed, termen sinuate, oblique; 4 and 5 coincident, 8 and 9 out of 7; light brownish irrorated dark fuscous and blackish; an irregular white streak from costa at $\frac{1}{3}$ to dorsum before middle, preceded by stronger blackish irroration, angulated below middle, above middle with blackish white-ringed dot resting on it (first discal stigma); second discal stigma transverse, of blackish irroration, ringed white; a rather thick irregular white streak from $\frac{3}{4}$ of costa to $\frac{4}{5}$ of dorsum, indented above middle, stronger blackish suffusion towards costa before and beyond this: cilia dark grey irrorated white, base white on termen. Hindwings grey, thinly scaled; cilia whitish, two grey shades.

S. INDIA, Ootacamund, April (*Mujtaba*); 1 ex.

Pachnistis solita, n. sp.

♂. 11 mm. Head, thorax light brownish-ochreous or whitish-ochreous, slightly speckled fuscous. Palpi ochreous-whitish, second joint irrorated dark fuscous except apex, anterior edge of terminal joint sprinkled dark fuscous. Forewings slightly dilated, apex obtuse, termen obliquely rounded; light brownish-ochreous or whitish-ochreous speckled fuscous; a fuscous or ferruginous dot near dorsum at base; stigmata small, dark fuscous or ferruginous, plical rather before first discal; an almost marginal series of dark fuscous or ferruginous dots round posterior part of costa and termen: cilia pale ochreous or ochreous-whitish, interrupted lines or faint bars of fuscous speckling. Hindwings grey, subdorsal hairs forming a long expansible ochreous-whitish tuft; cilia pale greyish or ochreous-whitish.

Fiji, Labasa, Lautoka, December to February, and bred from infested maize in July (*Veitch, Greenwood*); 6 ex. (type Brit. Mus.). This insect is so similar to *Stœberhinus testacea* that it would appear to be identical but for the difference in character of the ♂ palpi, and suggests interesting speculations.

Brachmia craterospila, n. sp.

♀. 13 mm. Head, thorax ochreous-whitish. Palpi ochreous-whitish, second joint dark fuscous except apex. Forewings termen slightly sinuate, oblique; 7 and 8 stalked, 7 to apex; ochreous-whitish, a few grey specks; a blackish-grey spot on base of costa, and a small spot on dorsum near base; stigmata large, blackish, plical slightly before first discal, second discal larger, forming a round spot; a small blackish spot on dorsum towards tornus; a præmarginal series of cloudy blackish dots round posterior part of costa and termen: cilia whitish-ochreous, some blackish specks. Hindwings light grey; cilia whitish, a greyish subbasal shade.

ASSAM, Shillong, September (*Fletcher*); 1 ex.

Brachmia xanthocosma, n. sp.

♂ ♀. 19-21 mm. Head orange. Palpi orange-yellow, base of second joint mixed dark grey, anterior edge of terminal joint dark fuscous. Antennæ yellow, apical fifth light grey. Thorax, abdomen dark purple-fuscous, venter light orange. Posterior legs orange-yellowish, femora and basal half of tibiæ dark fuscous. Forewings posteriorly dilated, termen slightly rounded, little oblique; 8 and 9 out of 7, 7 to apex; dark purple-fuscous: cilia concolorous. Hindwings and cilia dark fuscous.

UGANDA, Kampala (*Hargreaves*); 2 ex. (type Brit. Mus.).

Brachmia ptochodryas, n. sp.

♂. 12 mm. Head, thorax pale greyish-ochreous, thorax sprinkled fuscous. Palpi whitish, second joint sprinkled fuscous except apex, anterior edge of terminal joint sprinkled dark fuscous. Forewings apex obtuse-pointed, termen very obliquely rounded; 7 and 8 stalked, 7 to apex; light brownish-ochreous speckled fuscous; stigmata dark fuscous, plical little marked, somewhat before first discal, an undefined spot of dark fuscous suffusion on dorsal before second discal; a præmarginal series of cloudy dark fuscous dots round posterior part of costa and termen: cilia pale brownish-ochreous thinly speckled fuscous on basal half, paler towards tips. Hindwings grey; cilia light grey.

ASSAM, Shillong, 5000 feet, September (*Fletcher*); 2 ex.

Brachmia strophopa, n. sp.

♂. 21 mm. Head, thorax dark bronzy-fuscous. Palpi ♂ pale ochreous, ♀ ochreous-whitish, second joint fuscous externally except towards tip. Antennæ dark grey, in ♂ simple. Forewings somewhat dilated, termen slightly sinuate, rather oblique; 8 and 9 out of 7, 7 to just below apex; rather dark bronzy-fuscous; stigmata black, first discal moderate, others large, plical obliquely beyond first discal, second discal transverse; a hardly perceptible

lighter subterminal shade angulated outwards in middle and inwards near costa, on costa pale ochreous: cilia ♂ pale ochreous tinged grey, a rather dark fuscous subbasal shade, ♀ grey without darker shade. Hindwings light grey; cilia ♂ pale ochreous-greyish, a darker subbasal shade, ♀ greyer.

S. INDIA, Palnis, Kodaikanal, 7000 feet, August (*Fletcher*); 2 ex. Very like *rusticana*, but darker; in *rusticana* the antennæ in ♂ are rather strongly ciliated (not mentioned in description).

***Brachmia consummata*, n. sp.**

♂ ♀. 16–19 mm. Head, thorax light brownish-ochreous. Palpi pale yellow-ochreous more or less sprinkled fuscous. Forewings moderate, termen nearly straight, rather oblique; 7 and 8 stalked; ochreous, more or less sprinkled fuscous; stigmata black, plical beneath first discal; a small cloudy spot or short inwards-oblique streak of fuscous suffusion from dorsum towards tornus; a submarginal series of small irregular blackish dots round apex and termen: cilia light ochreous. Hindwings light ochreous-greyish; cilia pale ochreous, a pale greyish subbasal shade.

ASSAM, Shillong, 5000–6000 feet, June, July (*Fletcher*); 6 ex.

***Brachmia trichocyma*, n. sp.**

♂ ♀. 15 mm. Head grey or fuscous. Palpi fuscous, terminal joint lined white and black. Thorax fuscous, white supralateral streaks. Forewings termen nearly straight, scarcely oblique; 7 and 8 stalked, 7 to apex; rather dark fuscous; plical and first discal stigmata represented by an indistinct somewhat 8-shaped brownish spot edged with some blackish scales and then anteriorly by a fine white waved line extended to dorsum, second discal minute, blackish, followed by a white dot, beneath it a more or less developed oblique white strigula about fold; a small whitish spot on costa at $\frac{4}{5}$, whence a faint series of white scales crosses wing; indistinct small dark marginal dots round apex and termen, sometimes separated by small white marks: cilia grey. Hindwings dark grey; cilia light grey, a darker basal shade.

BRAZIL, R. Trombetas, Teffé, September, January; BRITISH GUIANA, Bartica, January (*Parish*); 4 ex. Allied to *convolvuli*.

***Brachmia cerinura*, n. sp.**

♂. 12–13 mm. Head, thorax fuscous. Palpi whitish-ochreous, second joint very stout, suffusedly mixed fuscous, terminal joint with two fuscous bands. Anal tuft yellow-ochreous. Forewings termen nearly straight, scarcely oblique; 7 and 8 stalked, 7 to apex; rather dark fuscous; plical and first discal stigmata represented by a rather 8-shaped brownish spot edged with some blackish scales, a short fine whitish line from dorsum preceding this, second discal forming a roundish similar spot; a small cloudy whitish

spot on costa at $\frac{5}{8}$; some marginal white specks round apex and termen; cilia grey, tips white on termen. Hindwings grey; cilia grey-whitish, a grey subbasal line.

BRAZIL, Obidos, September (*Parish*); 2 ex.

***Brachmia syntonopis*, n. sp.**

♂. 14 mm. Head light ochreous. Palpi pale ochreous, second joint blackish except apex, terminal joint with blackish subapical band. Thorax light ochreous, a spot on shoulder and two posterior spots blackish. Forewings termen rounded, rather strongly oblique; 7 and 8 stalked; light ochreous; markings blackish, partly with raised scales; a spot above base of dorsum; a slender streak along basal half of costa; an oblique fascia from base of costa to fold, terminating in two lobes representing plical and first discal stigmata, former somewhat anterior, a posterior projection from fascia towards costa; a rather large irregular spot on middle of costa; an excurved-crescentic spot from dorsum before tornus, touching a broad transverse fascia from costa at $\frac{3}{4}$ not reaching tornus; an irregular interrupted streak along termen; cilia yellow-ochreous, a faint interrupted median brownish shade. Hindwings grey; cilia light greyish, base whitish-ochreous.

BOMBAY, Belgaum, October (*Maxwell*); 1 ex.

***Glyphidocera percnoleuca*, n. sp.**

♂. 10 mm. Head, palpi, thorax fuscous. Antennal notch shallow. Forewings termen rounded, somewhat oblique; 7 and 9 stalked; pale fuscous irrorated dark fuscous; discal stigmata cloudy, dark fuscous; cilia whitish-grey, a dark grey shade. Hindwings and cilia ochreous-whitish.

BRAZIL, Parintins, October (*Parish*); 1 ex.

***Glyphidocera psammolitha*, n. sp.**

♂ ♀. 11-13 mm. Head, palpi, thorax pale greyish-ochreous sprinkled fuscous. Antennal notch ♂ slight, almost filled by scale-projection. Forewings termen rounded, little oblique; 7 and 9 usually stalked (9 examples stalked, 3 separate); light greyish-ochreous more or less irrorated fuscous or dark fuscous; stigmata forming small cloudy fuscous or dark fuscous spots or dots, plical rather obliquely before first discal, sometimes an additional spot midway between first discal and base; sometimes terminal dots of dark fuscous suffusion; cilia pale greyish-ochreous, two suffused grey shades. Hindwings ochreous-whitish, apex sometimes greyish-tinged; cilia ochreous-whitish.

BRAZIL, Santarem, Obidos, Tefé, August to January (*Parish*); 12 ex.

Glyphidocera inurbana Meyr.

Hindwings ♂ dorsal area clothed with rough hairscales (not noticed in my description). Occurs also at Parintins, Brazil, October.

Glyphidocera stenomorpha, n. sp.

♂. 13 mm. Head, thorax light yellow-ochreous. Palpi pale ochreous suffusedly irrorated dark fuscous. Antennal notch rather shallow, partly filled by scale-projection. Abdomen grey. Forewings rather narrow, termen rounded, rather oblique; 7 and 9 approximated at base; light yellow-ochreous, a few fuscous specks; stigmata forming small fuscous dots or marks, plical obliquely before first discal; a slender terminal streak of rather dark fuscous suffusion; cilia greyish mixed dark fuscous, on costa and beneath tornus light ochreous-yellowish. Hindwings $1\frac{1}{2}$, grey; cilia grey-whitish.

DUTCH GUIANA, Paramaribo, August; 1 ex.

Glyphidocera crocogramma, n. sp.

♂. 13 mm. Head pale ochreous, hairs of crown suffused fuscous. Palpi whitish-ochreous irrorated dark fuscous. Antennal notch rather slight, almost filled by scale-projection. Thorax pale ochreous, shoulders sprinkled fuscous. Forewings termen slightly rounded, somewhat oblique; 7 and 9 separate; pale ochreous sprinkled rather dark fuscous; discal stigmata forming small cloudy dark fuscous spots, plical less defined, somewhat before first discal: cilia grey-whitish, two grey shades. Hindwings whitish, tinged grey posteriorly; a light ochreous-yellow streak along vein 1 c from base to beyond middle (sexual); cilia whitish.

BRAZIL, R. Trombetas, September (*Parish*); 1 ex.

Glyphidocera orthoctenis, n. sp.

♂. 11 mm. Head, thorax pale yellow-ochreous, sometimes sprinkled fuscous. Palpi whitish-ochreous sprinkled dark fuscous. Antennal notch slight, filled by scale-projection. Forewings termen rounded, somewhat oblique; 7 and 9 separate; pale yellow-ochreous, more or less sprinkled fuscous or dark fuscous; discal stigmata rather small, dark fuscous, plical little marked: cilia grey, a dark grey subbasal shade. Hindwings light grey, suffused whitish-ochreous on anterior half; a fringe of long erect scales along vein 1 b; cilia ochreous-whitish.

BRAZIL, Para, July (*Parish*); 3 ex.

Glyphidocera catectis, n. sp.

♂. 12 mm. Head, thorax greyish-ochreous mixed fuscous. Palpi ochreous-whitish irrorated dark fuscous. Antennal notch deep, forming with strong projection a circular excavation. Abdomen ochreous-whitish. Forewings termen nearly straight, little oblique;

7 and 9 separate; ochreous irrorated fuscous; a rather large transverse-oval fuscous spot in disc at $\frac{2}{5}$, and a smaller transverse spot at $\frac{2}{3}$, space between these clearer ochreous: cilia ochreous-whitish, on costa more ochreous, at apex tinged grey, on termen basal third irrorated fuscous. Hindwings pale grey, basal half suffused ochreous-whitish and partially clothed with modified scales (sexual); cilia ochreous-whitish.

ECUADOR, Huigra, 4500 feet, June (*Parish*); 2 ex.

Autosticha stigmatopis, n. sp.

♀. 14-16 mm. Head, thorax light fuscous, somewhat sprinkled whitish, in one example head mostly suffused whitish. Palpi grey speckled whitish, apex of second joint whitish. Forewings elongate, apex obtuse, termen very obliquely rounded; light fuscous somewhat whitish-speckled; stigmata large, cloudy, dark fuscous, plical beneath first discal; a præmarginal series of cloudy groups of dark fuscous scales round posterior part of costa and termen to beneath second discal: cilia pale grey, base sprinkled darker. Hindwings grey; cilia pale grey.

S. INDIA, Cuddapah, 4000 feet (*Campbell*); 5 ex.

Autosticha thermopis, n. sp.

♂. 15 mm. Head, thorax ferruginous-brown. Palpi ferruginous, terminal joint dark grey, ochreous-whitish on sides towards base. Antennal ciliations 3, fasciculated. Forewings moderate, termen hardly rounded, oblique; ferruginous-brownish; stigmata large, cloudy, darker, plical beneath first discal, second discal transverse; a præmarginal series of faint cloudy similar spots round posterior part of costa and termen: cilia light ferruginous-brownish. Hindwings dark grey; cilia light grey-brownish.

CEYLON, Maskeliya, December (*Pole*); 1 ex.

Encrasima xanthoclista, n. sp.

♂. 17 mm. Head ochreous-yellow. Palpi whitish-ochreous, second joint fuscous except tip. Thorax fuscous, shoulders ochreous-yellow. Forewings moderate, termen hardly rounded, rather oblique; grey-brownish; costa slenderly ochreous-yellow from base to near apex; stigmata moderate, black, plical slightly before first discal; a præmarginal series of cloudy blackish dots round posterior part of costa and termen, on termen obscured in a streak of grey suffusion: cilia ochreous-yellow, on tornus light grey, at apex a bar of grey suffusion. Hindwings grey; cilia light greyish, on upper part of termen tinged pale yellowish.

CEYLON, Madulsima, March (*Green*); 1 ex.

Phthorimaea extorris, n. sp.

♂ ♀. 11-12 mm. Head, thorax ochreous-grey-whitish speckled dark fuscous, face whitish. Palpi whitish, second and terminal

joints each with two bands of blackish irroration. Forewings 6 separate; whitish or ochreous-whitish closely irrorated dark grey; indistinct small dark grey spots on fold at base and $\frac{1}{5}$, and on costa between these; an oblique streak obscurely indicated by a dark grey spot on costa at $\frac{1}{4}$ and suffusion on fold beyond it; stigmata dark fuscous, sometimes distinctly ringed ochreous-brownish or whitish, plical beneath first discal, an additional dot midway between and above first and second discal; an angulated whitish subterminal line sometimes distinct in ♀, usually nearly obsolete, sometimes brownish-streaked on veins: cilia grey-whitish or ochreous-whitish, scattered blackish points. Hindwings grey, darker towards costa, lighter and more thinly scaled on median longitudinal third; cilia whitish-grey.

EGYPT, Aswan, March, April (*Hayward*); 8 ex.

***Phthorimaea crocoleuca*, n. sp.**

♀. 8 mm. Head, thorax white. Palpi white, second and terminal joints each with two rings of dark grey irroration. Forewings whitish; four moderately broad undefined irregular light ochreous-yellowish transverse fasciæ, costal edge slightly speckled blackish-grey on these, first almost basal, fourth almost apical: cilia white, some scattered black specks, base ochreous-yellowish round apex. Hindwings and cilia whitish.

EGYPT, Aswan, October (*Hayward*); 1 ex.

***Apatetris albiramis*, n. sp.**

♂. 9 mm. Head white. Palpi white, second joint with a grey lateral streak. Thorax whitish, four pale ochreous stripes. Forewings lanceolate, apex acutely produced; 7 and 8 out of 6; light yellow-ochreous; irregular white streaks along all margins of cell and veins 7-11, partially sprinkled dark fuscous scales on edges, a slender similar streak along dorsum and termen throughout; an undefined streak of white suffusion along vein 1b: cilia white, a median line of fuscous points. Hindwings emargination obtuse, produced apex $\frac{1}{5}$; grey-whitish; cilia yellow-whitish, yellower towards base.

EGYPT, Aswan, April (*Hayward*); 1 ex.

METACHANDIDÆ.

***Ancylometis asbolopa*, n. sp.**

♂. 10 mm. Head, thorax dark grey. Palpi dark grey, tip of second joint whitish with black subapical ring, terminal joint whitish with two blackish bands. Forewings elongate, apex obtuse-pointed, termen very obliquely rounded; dark grey; an indistinct dark fuscous transverse streak before middle, bearing black tufts near costa and dorsum, their posterior edge dark grey speckled

whitish; two small black spots transversely placed on end of cell; a slender angulated fascia of blackish suffusion at $\frac{3}{4}$, beyond this a costal blotch of whitish speckling; an apical blotch of blackish suffusion: cilia grey. Hindwings dark grey; cilia grey.

S. INDIA, Palnis, Kodaikanal, 7000 feet, September (*Fletcher*); 1 ex.

EPERMENIADÆ.

Epermenia symmorias, n. sp.

♀. 10 mm. Head whitish-grey, crown speckled grey. Palpi whitish irrorated dark fuscous. Forewings rather narrow; grey-whitish speckled grey; a black dot beneath fold at $\frac{1}{5}$; a narrow rather dark brown antemedian fascia rather inwards-oblique from costa, dark fuscous towards costa, beyond this costal third of wing dark fuscous, median third rather dark brown, dorsal third irrorated dark grey with a projection into brown area at $\frac{3}{4}$ including a black dot; a fine oblique whitish strigula from costa towards apex; a fine blackish dash before apex: cilia round apex brown, a white basal line on costa, a black median line edged anteriorly on costa with white, beyond this a black apical spot edged above and below by white spaces, on tornus wholly grey. Hindwings and cilia grey.

FIJI, Labasa, November, bred from larva feeding on flowers of an unidentified plant (*Greenwood*); 1 ex. (Brit. Mus.).

ADELIDÆ.

Nemotois argyrospila, n. sp.

♂. 15 mm. Head black, hairs of back of crown light brownish-ochreous, eyes large, nearly approximated above. Palpi with very long hairs, externally blackish, internally whitish. Antennæ white, slight roughened black scales above near base. Thorax blackish. Forewings elongate, termen very obliquely rounded; shining golden-bronze; some undefined silvery-white suffusion on transverse vein: cilia grey, basal third coppery. Hindwings grey-whitish; cilia white, a blackish-grey basal line, above apex a grey patch.

PALESTINE, Haifa, March (*Buxton*); 1 ex. (Brit. Mus.).

PHALONIADÆ.

Phalonia exaequata, n. sp.

♂. 9 mm. Head ochreous-white, sides of crown tinged ferruginous. Palpi pale ferruginous-ochreous, tip white. Thorax whitish, shoulders ferruginous. Forewings slightly dilated, termen

rather obliquely rounded; ochreous-whitish; markings ferruginous suffusedly irrorated dark grey; a thick suffused streak along basal fourth of costa, and a suffused spot on dorsum near base; a moderate hardly curved fascia from beyond middle of costa to beyond middle of dorsum; one or two strigulæ on costa and dorsum beyond this; an irregularly triangular spot on dorsum before termen; a moderate fascia from costa near apex to lower half of termen, and some irregular marking on apical edge: cilia whitish-ochreous, a dark grey subbasal line. Hindwings grey; cilia grey-whitish, a grey subbasal shade.

PALESTINE, Haifa, May (*Buxton*); 1 ex. (Brit. Mus.).

Phalonia reclusa, n. sp.

♀. 13 mm. Head, palpi, thorax, abdomen white. Forewings termen hardly rounded, oblique; white; two rather narrow irregular ferruginous-ochreous fasciæ parallel to termen, interrupted beneath costa, first before middle, second towards termen: cilia white. Hindwings white, tinged ochreous posteriorly; cilia white.

PALESTINE, Tabagha, Galilee, April (*Buxton*); 1 ex. (Brit. Mus.).

LAMPRONIADÆ.

Chalceopla evocata, n. sp.

♂. 9 mm. Head, palpi pale ochreous. Thorax purple-grey. Forewings rather elongate, dilated, costa moderately arched, apex obtuse, termen obliquely rounded; 7-9 approximated at base; deep greyish-purple: cilia rather dark grey. Hindwings greyish-purple; cilia grey.

ASSAM, Cherrapunji (*Fletcher*); 2 ex.

TORTRICIDÆ.

Cacoecia euryplintha, n. sp.

♀. 20 mm. Head, palpi, thorax ferruginous-brownish. Forewings broad-oblong, costa anteriorly very strongly arched, posteriorly slightly sinuate, apex obtuse-prominent, termen rather sinuate, vertical; purplish-ochreous-brown, scattered transverse deep ferruginous strigæ; basal patch with rounded edge, and oblique central fascia indicated by rather darker suffusion, edged similar striæ; undefined elongate costal patch indicated by dark ferruginous suffusion. Hindwings orange, dorsal half dark grey; costal cilia forming a thickened grey patch from $\frac{2}{5}$ to near apex.

BENGAL, Darjiling, Mangpu, 4000 feet, September (*Fletcher*); 1 ex.

Cacoecia mansueta, n. sp.

♂. 24 mm. Head, thorax pale greyish-ochreous, two spots on face, collar, edge of shoulder, and palpi dark reddish-fuscous. Antennal ciliations fasciculated (2). Forewings oblong, costa anteriorly strongly arched, without fold, posteriorly sinuate, apex obtuse, termen almost straight, little oblique; brownish-ochreous, costal half largely irregularly suffused lilac-fuscous, darkest along costa; basal patch outlined by a faint pale sinuate transverse stria; two short similar striae crossing dorsal area beyond this; central fascia outlined by two similar striae, rather narrow, somewhat oblique, excurved on lower $\frac{2}{3}$; two similar striae from costa towards apex to tornus: cilia light greyish-fuscous, becoming fuscous round apex (imperfect). Hindwings ochreous-grey-whitish, suffused pale grey towards apex; cilia ochreous-whitish.

BRAZIL, Rio Janeiro; 1 ex. (Coll. Joannis).

Cacoecia interjecta Meyr.

Locality (accidentally omitted) BENGAL, Darjiling; 1 ex. (Coll. Joannis).

Cacoecia expleta, n. sp.

♂. 21-22 mm., ♀ 27-28 mm. Head, palpi, thorax brown. Forewings suboblong, costa ♂ anteriorly strongly arched, edge very slightly folded on basal half with short scale-tooth towards base, posteriorly faintly sinuate, ♀ more strongly arched and sinuate, apex obtuse, termen vertical, ♂ hardly sinuate, ♀ more sinuate; light brown, with fine irregular interrupted leaden-grey transverse strigæ and lines on veins; markings darker brown, edged dark grey strigæ; basal patch hardly darker, edge sub-sinuate, oblique; central fascia moderate, oblique, posterior edge undefined except towards costa; costal patch semioval, ♀ larger and with a fasciate lobe posteriorly extending nearly to termen: cilia dark grey, tips light brownish. Hindwings orange, posteriorly with some faint greyish-tinged fine transverse strigulæ; cilia whitish-yellowish, an orange basal line.

ASSAM, Shillong, May, October (*Fletcher*); 4 ex.

Tortrix velitans, n. sp.

♂. 15 mm. Head, thorax whity-brownish. (Palpi missing.) Antennæ serrulate, minutely ciliated. Forewings moderate, costa anteriorly moderately arched, with narrow fold from base to $\frac{2}{5}$, then nearly straight, termen rounded, somewhat oblique; whitish; costal fold dark reddish-fuscous; edge of basal patch indicated by a blackish spot on fold connected with costa by a very fine blackish curved line enlarged into a small mark above middle; some crimson and grey irroration towards dorsum beneath and beyond this; central fascia moderate, rather oblique, upper part greyish, dark

grey on costa, edged dark reddish-fuscous beneath costa, in disc crossed by a large dark reddish-fuscous spot, nearly interrupted above this, below it indicated by some brownish suffusion, towards dorsum by crimson-fuscous irroration only; costal patch elongate-semioval, reddish-grey, darker towards costa; some light greyish suffusion along termen, with fine strigulæ of blackish irroration: cilia grey, on apex and tornus whitish. Hindwings ochreous-whitish; a few grey strigulæ towards apex; cilia ochreous-whitish.

BRAZIL, probably south-east district; 1 ex. (Coll. Joannis).

Tortrix liobathra, n. sp.

♂. 13 mm. Head, palpi, thorax brownish. Forewings sub-oblong, costa bent before middle, with moderate fold from base to near middle, apex obtuse-pointed, termen nearly straight, oblique; fuscous, from base to central fascia smooth, uniform, fold brownish-tinged, on dorsum an elongate-semioval patch of modified brownish-ochreous scales; central fascia narrow, little oblique, dark ferruginous-brown mixed blackish, somewhat dilated towards dorsum, edged anteriorly by a light brownish stria; groundcolour beyond this darker purplish-fuscous, much mixed and suffused ferruginous-brown, costal patch semioval, darker, some strigulæ of blackish scales round this and before termen: cilia light ochreous, an interrupted dark grey subbasal shade. Hindwings rather dark grey, thinly scaled anteriorly; cilia grey.

ASSAM, Shillong, September (*Fletcher*); 1 ex.

DELTOBATHRA, n. g.

Antennæ ♂ minutely ciliated. Palpi moderate, curved, ascending, with appressed scales, terminal joint short. Thorax without crest. Forewings 3 from angle, 7 to termen. Hindwings without cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 stalked.

Type *platamodes* Meyr. I refer here also *nigrivelata* Wals., described as a *Tortrix*, of which I have a long series from the Amazons.

Deltobathra platamodes, n. sp.

♂ ♀. 11-12 mm. Head, palpi, thorax dark fuscous. Forewings elongate, dilated, costa gently arched, termen rounded, somewhat oblique; brownish-ochreous; a small triangular dark brown or dark fuscous basal patch, very narrow on dorsum; central fascia rather broad, oblique, brown or dark brown, posteriorly edged blackish scales; wing beyond this mostly suffused pale iridescent-grey; costal spot subquadrate, ferruginous-brown, edged with some blackish scales, beneath it a roundish ferruginous-brown spot irrorated or irregularly marked blackish resting on middle of termen; cilia iridescent brownish-ochreous, on middle of

termen two or three indistinct blackish bars. Hindwings dark grey; cilia light grey.

BRAZIL, Para, July; PERU, Jurimaguas, March (*Parish*); 3 ex.

Peronea compsoptila, n. sp.

♀. 19 mm. Head grey-whitish somewhat mixed grey. Palpi grey sprinkled whitish. Thorax lilac-brown. Forewings elongate, costa strongly arched, apex obtuse-pointed, termen slightly sinuate, somewhat oblique; violet-brown, lighter and ochreous-suffused towards costa beyond middle, apical area darker red-brown, on dorsal half of wing veins marked whitish lines and interspaces more or less suffused whitish irroration; several small blackish tufts accompanied with a few white scales in a series just below fold, one above fold in middle, an oblique series of four from beneath costa before middle to below middle of disc, and two transversely placed in disc towards apex; two or three snow-white dots and scattered scales at apex: cilia dark grey speckled whitish, basal third suffused pale brownish, beneath tornus ochreous-whitish. Hindwings grey, lighter anteriorly; cilia whitish, a grey subbasal line.

ASSAM, Shillong, November (*Fletcher*); 1 ex.

COSMOPTERYGIDÆ.

Labdia clytemnestra, n. sp.

♂ ♀. 11–12 mm. Head rosy-grey-whitish. Palpi stouter than in *semicoccinea*, ochreous-whitish, terminal joint with incomplete basal and submedian blackish rings and sharp subapical band. Antennæ whitish ringed blackish, a blackish band at $\frac{3}{4}$, above this whitish with two slender blackish bands. Thorax reddish-orange, four pale greyish stripes. Abdomen greyish, two basal segments pale ochreous. Posterior tibiæ above with orange streak and whitish-ochreous hairs. Forewings basal (larger) half vermilion-red, four blue-grey streaks, first zigzag from base of costa to near dorsum about $\frac{1}{4}$, second zigzag from costa near base obliquely to fold, thence along fold to near middle and bent up to meet apex of third, third from or near fourth near origin above middle to edge of red area, fourth costal from $\frac{1}{5}$ to middle, its extreme costal edge whitish; a grey-whitish dorsal streak speckled blackish from near base to near edge of red area, connected by slight whitish-grey bars with second grey streak at its middle and near extremity; posterior area pale ochreous, a fine dash projecting into third grey streak, on dorsal half a pale golden-metallic bar separating the colours with a slender discal projection to tornus and followed in disc by a minute black dot, on costa minute black dots surrounded whitish just beyond red area and at $\frac{2}{3}$: cilia pale ochreous. Hindwings and cilia light grey, cilia above apex suffused ochreous-whitish. Forewings ♂

beneath with thick subdorsal streak of dense raised greyish-fulvous scales extending from vein 2 halfway to apex.

FIJI, Lautoka, June, July, bred from dead branches of *Mangifera* (*Greenwood, Veitch*); 3 ex. (type Brit. Mus.). The blackish specks in dorsal streak distinguish this from any of the similar species.

***Labdia scenodoxa*, n. sp.**

♂. 11–12 mm. Head pale ochreous, face white, eyes crimson. Palpi ochreous, anterior edge of terminal joint and a subapical ring blackish, tip white. Thorax pale silvery-grey, an orange-red longitudinal mark anteriorly on each side of back (sometimes obsolete?), an expansible pencil of long whitish hairs lying beneath forewings. Abdomen silvery-whitish. Forewings basal (larger) half orange-red, five light grey-violet-blue streaks, first from base along dorsum to middle of wing, second from base of fold running into first before extremity, third from base of costa rather irregular and thence along fold and turned up at apex to join apex of fourth, fourth out of fifth near anterior extremity, then above middle to end of red area, fifth along costa from $\frac{1}{5}$ to end of red area; posterior area light pinkish-ochreous, at anterior edge a black whitish-circled costal dot and a silvery dorsal spot surmounted by a black dot, a small black mark on costa at $\frac{2}{3}$ edged anteriorly white and posteriorly by a wedgeshaped spot of white suffusion, beyond this at tornus a small spot of silvery-whitish suffusion sometimes accompanied by one or two black scales: cilia pale bronzy-ochreous, towards tornus greyish. Hindwings and cilia light grey.

FIJI, Labasa, September (*Veitch*); 2 ex. (type Brit. Mus.).

***Labdia cosmangela*, n. sp.**

♂ ♀. 14–15 mm. Head whitish, eyes crimson. Palpi very slender, whitish, terminal joint with indistinct greyish submedian and subapical bands. Antennæ whitish ringed blackish-grey, apical third whitish with two grey bands. Thorax orange, four pale grey stripes. Abdomen pale ochreous. Posterior tibiæ above orange-yellowish with ochreous-whitish hairs. Forewings basal $\frac{3}{5}$ deep orange, four whitish-blue-grey streaks, first along fold from base almost to edge of orange area, turned up at tip to meet third, second rather irregular from costa near base to middle of first, third from above end of second to edge of orange area, fourth costal from $\frac{1}{5}$ to edge of orange area, its extreme costal edge whitish; an irregular grey-whitish dorsal streak from base to near edge of orange area, connected by some whitish suffusion with plical streak at $\frac{1}{3}$ of wing; posterior area pale ochreous, a fine dash projecting into third grey streak, a silver-metallic mark surmounted by a blackish dot on dorsum separating the two colours and a small silver-metallic spot on termen nearer to this than to apex, on costa short blackish transverse marks surrounded white just beyond orange area and at $\frac{2}{3}$: cilia pale ochreous. Hindwings rather

dark grey; cilia pale greyish, above apex ochreous-whitish, towards dorsum suffused whitish-ochreous. Forewings ♂ beneath with small greyish-fulvous glandular dorsal spot between veins 3 and 4.

QUEENSLAND, Cairns, September to November (*Dodd*); 3 ex. In *semicoccinea* Staint. (with which I had hitherto erroneously confounded this species and *clytemnestra*) the ♂ has a long expansible pencil of ochreous-yellowish hairs from metathorax lying along each side of abdomen; in the allied *molybdaula* the ♂ has a long expansible pencil of fine whitish hairs projecting from mesothorax above hindwings; neither of these two species has yet been found in the Australian region.

Labdia microdictyas, n. sp.

♀. 7 mm. Head, thorax white. Palpi white, subapical ring of second joint and submedian and subapical rings of terminal joint grey. Abdomen whitish, two basal segments pale ochreous. Forewings anterior $\frac{2}{5}$ reddish-orange, with iridescent pale violet-grey markings, viz., a broad irregular streak along dorsum to middle, at base extending to costa, an oblique streak from costa at $\frac{1}{5}$ running into this, and an oblique very irregular streak from costa before middle to edge of orange area, where it is slenderly connected with apex of dorsal streak; orange area limited by a slender silvery-whitish somewhat oblique streak edged posteriorly with some black scales except in middle: posterior area yellow-ochreous, including an undefined blotch of blue-whitish suffusion resting on costa, preceded on costa and followed on tornus by small dots of black irroration: cilia whitish. Hindwings pale grey; cilia ochreous-whitish.

FILE, Labasa, September (*Veitch*); 1 ex. (Brit. Mus.).

Labdia allotriopa, n. sp.

♂ ♀. 8-9 mm. Head ♂ white, ♀ grey-whitish. Palpi white, second joint dark fuscous except apex, terminal joint infuscated towards apex. Thorax ♂ white, ♀ grey-whitish, patagia dark brown-grey, sometimes a dark central line. Forewings dark brown-grey; an irregular white dorsal stripe from base to tornus, in ♀ greyish-tinged, irregularly expanded beyond middle, including an elongate blackish mark on dorsum in middle; in ♂ a more or less developed irregular streak of white suffusion along costa from near base to near apex, interrupted in middle, in ♀ reduced to a posterior spot, a black mark beneath this just before apex; a rather large black dot partly white-circled, resting on edge of dorsal streak beyond middle of wing, and a similar dot in disc near beyond it (second discal stigma); along termen a series of several dark dots surrounded white and edged anteriorly by a fine white streak: cilia light grey, towards tips whitish, round apex a dark grey sabbasal

shade. Hindwings and cilia grey; in ♂ a long expansible hair-pencil from base lying along costa.

FIJI, Kabulu, bred October from ripe cotton-bolls (*Simmonds*), Labasa, August (*Greenwood*); 4 ex.

***Labdia intuens*, n. sp.**

♂. 10 mm. (Head missing.) Thorax fuscous, two triangular white dorsal spots nearly meeting. Forewings rather broad-lanceolate: brown; a blackish dash beneath fold from base to $\frac{1}{4}$, edged whitish suffusion beneath; a short very oblique blackish streak from costa near base, from its apex a subcostal streak of whitish irroration to near middle; short very oblique blackish marks from middle of costa and middle of dorsum, former followed by a white costal spot; plical and second discal stigmata large, approximated, black, irregularly circled white, second discal connected by white suffusion with median costal spot and with a smaller white spot on costa towards apex, followed by blackish suffusion extending to apex; a white terminal line: cilia grey, round apex white within a blackish antemedian line, beneath apex base more broadly white. Hindwings dark grey; cilia grey.

FIJI, Lautoka, July (*Greenwood*); 1 ex. (Brit. Mus.).

***Pyroderces megacentra*, n. sp.**

♀. 12 mm. Head rosy-whitish, slightly sprinkled light grey. Palpi grey-whitish, second and terminal joints each with three blackish rings. Thorax pale pinkish, an antemedian transverse band of dark grey irroration. Forewings pale pinkish; a dark grey basal patch occupying $\frac{1}{3}$ of wing, including a spot of ground-colour below fold; separated from this by a narrow space, in middle of which is some dark grey suffusion, is a rather broad dark grey postmedian fascia, outwards-oblique from dorsum, on middle of its anterior edge a large blackish dot white-circled above (first discal stigma); a rather broad dark grey streak along termen, tending to be connected with preceding by three or four narrow bars of dark grey irroration, on middle of anterior edge of this a large black dot (second discal stigma), on costal preceded by a slight white mark: cilia light grey, round apex pale pinkish, at apex a short blackish basal bar with projections at extremities, before apex a small blackish-grey spot. Hindwings dark grey; cilia grey.

FIJI, Dreketi R., September, at light (*Simmonds*); 1 ex. (Brit. Mus.).

***Limnoecia inconcinna*, n. sp.**

♂. 9 mm. Head, thorax purplish-grey. Palpi white, anteriorly dark grey. Antennæ black finely lined white. Forewings dark purplish-grey sprinkled whitish; plical and second discal stigmata blackish, obscurely whitish-ringed; a small obscure whitish spot on costa beyond middle; a more distinct whitish oblique wedge-shaped mark from costa towards apex, partly in cilia; some white

marginal specks round apex and termen: cilia grey, at apex a darker subbasal shade. Hindwings grey; an ochreous-whitish hairpencil from base lying along costa; cilia grey.

FIJI, Labasa, Kabulu, August (*Greenwood*): 3 ex. (type Brit. Mus.).

Persicoptila phoenoxantha, n. sp.

♀. 12–13 mm. Head, palpi, thorax ochreous-yellowish. Posterior tibiæ rose-pink barred dark grey. Forewings crimson mixed or suffused dark grey; base ochreous-yellow; a moderately broad somewhat oblique ochreous-yellow fascia beyond middle narrowed towards dorsum; apex orange-yellow: cilia round apex pale orange, towards tornus light grey. Hindwings grey; cilia light grey.

FIJI, Labasa, September, October (*Veitch*); 2 ex. (type Brit. Mus.).

Ascalenia armigera, n. sp.

♂. 9 mm. Head, thorax dark fuscous. Palpi dark fuscous slightly sprinkled whitish. Forewings whitish-ochreous; base rather narrowly dark fuscous, edge blackish; a moderate streak of rather dark fuscous suffusion along costa from $\frac{1}{4}$ to apex, including a white dot at $\frac{3}{4}$; stigmata moderate, black, plical rather before first discal; slight fuscous suffusion towards dorsum posteriorly: cilia rather dark fuscous. Hindwings dark grey; cilia greyish.

FIJI, Labasa, November (*Greenwood*); 1 ex. (Brit. Mus.).

Ascalenia subcrocea, n. sp.

♂. 7 mm. Head, thorax greyish irrorated darker, face ochreous-whitish. Palpi ochreous-whitish, terminal joint speckled blackish anteriorly. Forewings greyish, tips of scales dark grey, forming a fine transverse striation; stigmata large, blackish-grey, plical beneath first discal; small cloudy grey-whitish opposite costal and tornal spots at $\frac{3}{4}$, apical area beyond this darker-suffused: cilia ochreous-grey-whitish, base mixed dark grey. Hindwings grey-whitish, apex yellowish-tinged; cilia whitish-yellowish.

EGYPT, Aswan, March (*Hayward*); 1 ex.

ELACHISTIDÆ.

Elachista putris, n. sp.

♂. 8 mm. Head grey, face whitish. Palpi dark grey, extreme tip of second joint whitish. Thorax dark grey. Forewings lanceolate; dark purplish-grey irrorated blackish, posteriorly slightly whitish-speckled; a narrow irregular-edged slightly oblique ochreous-white fascia before middle, on costa narrower and with some blackish scales; an irregular-edged ochreous-white spot with some blackish scales on costa at $\frac{3}{4}$, and a small similar spot or dot on dorsum wholly before it: cilia grey, within a blackish postmedian line irrorated blackish and speckled white. Hindwings dark grey; cilia grey.

PUNJAB, Murree, 7500 feet, May (*Dutt*); 2 ex.

Elachista leucobathra, n. sp.

♂. 8 mm. Head ochreous-whitish irrorated blackish, on face more sparsely. Palpi dark grey, tip of second joint whitish. Thorax blackish, apex of patagia mixed whitish. Forewings lanceolate, acute; blackish; a narrow white basal fascia, edge rather oblique; a narrow irregular-edged almost direct silvery-white fascia before middle; a subtriangular silvery-white spot on costa at $\frac{3}{4}$, and a dot or small spot on tornus wholly before it: cilia grey, a black median shade. Hindwings dark grey; cilia grey.

PUNJAB, Murree, 7500 feet, May (*Dutt*); 2 ex.

OECOPHORIDÆ.

Promalactis commotica, n. sp.

♂ ♀. 9 mm. Head white, sidetufts bronzy-yellow, face fuscous. Palpi second joint dark grey, terminal white with blackish median band. Thorax bronzy-yellow. Forewings elongate-lanceolate; deep ochreous-yellow; three slender white streaks from dorsum edged blackish scales, first almost along basal portion, second from $\frac{1}{3}$, erect, reaching more than half across wing, third from $\frac{2}{3}$, oblique, reaching half across wing; a rounded-transverse white spot from costa beyond middle reaching half across wing, preceded by some slight dark grey suffusion towards costa; a small obscure white spot on tornus, preceded and followed by blackish scales along margin; an irregular-triangular white apical spot edged black irroration except on costa: cilia ochreous-yellowish, some black speckling towards tornus. Hindwings grey; cilia light grey.

ASSAM, Shillong, July (*Fletcher*); 2 ex.

Nochelodes temperatella Led.

This specific name supersedes *xenicopa* Meyr.; in Staudinger's Catalogue the species is referred by Rebel to *Scythris*, which led to my overlooking it, as it has unquestionably no affinity with that genus.

Pleurota platyrrhoa, n. sp.

♂. 24 mm. Head, thorax white. (Palpi missing.) Forewings apex pointed, termen very obliquely rounded; light yellowish-ochreous; costal edge obscurely whitish from near base to about middle; a broad snow-white median stripe from base to apex, rather narrowed in middle, from before this sending a broad rather uneven branch along fold to extremity; on anterior half of dorsum some broad undefined white suffusion: cilia ochreous-whitish. Hindwings grey, towards apex and upper part of termen whitish-mixed; cilia ochreous-whitish, a faint light greyish subbasal shade.

PALESTINE, Jerusalem, April (*Buxton*); 1 ex. (Brit. Mus.). Nearest *oranella*.

Pleurota idalia, n. sp.

♂. 15-18 mm. Head, thorax whitish, shoulders fuscous, eyes dark reddish. Palpi grey, upper hairs of second joint whitish, terminal joint not much shorter than second, whitish posteriorly. Forewings apex pointed, termen very obliquely rounded; white, hardly perceptibly ochreous tinged; a moderately broad fuscous streak from base of costa to costa just above apex, leaving costa narrowly white between its ends: cilia white, on costal end of streak light greyish. Hindwings rather dark grey; cilia whitish-grey.

CYPRUS, Forest St. Nicolas, May (*Mavromoustakis*); 6 ex. (type Brit. Mus.).

Peritorneuta gennaea, n. sp.

♂. 15 mm. Head, thorax whitish-ochreous-grey. Palpi grey, tip whitish. Forewings whitish closely irrorated grey; scattered small black dots, especially subcostal and subdorsal series posteriorly, and on veins towards termen, veins white-lined near termen; a short streak of pinkish suffusion on fold beneath middle of wing; a subrescentic blackish mark on end of cell, beyond this a cloudy blotch of darker grey irroration suffused pinkish; a suffused darker grey semioval blotch on costa somewhat beyond middle; a curved-angulated whitish shade at $\frac{3}{4}$, followed by some darker grey suffusion with slight pinkish tinge: cilia whitish, a faint greyish subbasal line. Hindwings grey; cilia as in forewings.

FIJI, Labasa, September, bred from an unknown plant, larva short, broad, red and yellow (*Greenwood*); 1 ex. (Brit. Mus.).

Periacma eporphna, n. sp.

♂ ♀. 9-10 mm. Head light ochreous. Palpi pale ochreous, second joint with suffused dark fuscous subapical ring, terminal joint anteriorly more or less irrorated dark fuscous, in ♂ present but very short. Thorax ochreous more or less suffused purplish-grey. Forewings apex tolerably pointed, termen very obliquely rounded; 7 to costa; fuscous obscurely and suffusedly irrorated light ochreous; stigmata dark fuscous, plical slightly before first discal, sometimes obsolete, second discal transverse; blotches of light ochreous suffusion on dorsum before tornus, and on costa at middle and $\frac{3}{4}$, between these dark fuscous costal and tornal blotches connected by second discal stigma, and confluent beneath with a rather broad dark fuscous terminal fascia: cilia grey, at apex a patch of pale ochreous suffusion, a darker subbasal line. Hindwings dark grey; cilia grey.

ASSAM, Shillong, June, July (*Fletcher*); 5 ex. The structure of palpi is somewhat peculiar, but the natural affinity is here.

Pseudodoxia miseranda, n. sp.

♀. 20 mm. Head grey, sides of crown mixed ochreous-whitish. Palpi dark grey irregularly mixed whitish. Thorax grey. Forewings

apex obtuse, termen rounded, rather strongly oblique; light grey, irregularly mixed dark fuscous; stigmata moderately large, dark fuscous, plical beneath first discal, a streak of ochreous-whitish suffusion joining first and second discal; an acutely angulated almost marginal series of cloudy dots of dark fuscous irroration round posterior part of costa and termen: cilia whitish-grey, a broad grey antemedian shade. Hindwings grey, lighter between veins; cilia whitish-grey, a darker subbasal shade.

S. INDIA, Palnis, Kodaikanal, 7000 feet, August (*Fletcher*); 1 ex.

Hypersymmoca psammocryptis, n. sp.

♂. 13–15 mm. Head ochreous-whitish, slightly sprinkled dark fuscous. Palpi whitish, irrorated dark fuscous except apex of joints. Antennæ serrulate, with moderate ciliations (1) and pairs of longer single cilia (3) at apex of each joint. Thorax ochreous-whitish, more or less irrorated dark fuscous. Forewings elongate, apex obtuse, termen very obliquely rounded; ochreous-whitish variably irrorated dark fuscous, sometimes uniform, sometimes irregular or tending to form broad longitudinal streaks, sometimes with clear pale median and subdorsal streaks; second discal stigma forming a round usually distinct spot of dark fuscous irroration, others similar but usually more or less obscure or sometimes obsolete, plical beneath first discal: cilia ochreous-whitish, basal half sprinkled dark fuscous. Hindwings and cilia ochreous-grey-whitish.

EGYPT, Aswan, March, April (*Hayward*); 11 ex. Varies considerably, and should be near *facivorella* (which I have not seen), but the different ciliations of ♂ antennæ are distinctive.

LYONETIADÆ.

Hieroxestis divulsa, n. sp.

♀. 9–10 mm. Head ochreous-grey-whitish, with large fuscous frontal tuft extending below antennæ. Palpi pale greyish-ochreous, terminal joint fuscous, tip whitish. Thorax ochreous-grey-whitish, shoulders dark fuscous. Forewings lanceolate, acute; whitish-grey-ochreous, a few scattered long dark fuscous scales; a broad suffused dark fuscous streak from base (where it extends along basal portion of costa) above fold throughout and continued along termen to apex: cilia whitish-grey-ochreous, base mixed fuscous on termen. Hindwings light grey; cilia grey-whitish.

EGYPT, Aswan, March, April (*Hayward*); 4 ex.

PHRURIASTIS, n. g.

Head shortly rough-haired; ocelli posterior; tongue obsolete. Antennæ $\frac{4}{5}$, in ♂ simple, basal joint short, without pecten. Labial palpi moderately long, curved, ascending, rather stout, smooth,

terminal joint shorter than second, compressed, somewhat pointed. Maxillary palpi several-jointed, folded, slender, filiform. Posterior tibiæ smooth. Forewings apex somewhat down-turned; 2 from towards angle, 3 absent, 6 and 7 stalked, 7 to costa, 11 from middle. Hindwings $\frac{2}{3}$, lanceolate, cilia $1\frac{1}{2}$; cell open between 2 and 5, 3 and 4 absent, 6 absent, 5 and 7 approximated.

Belongs to group of *Erechthias*.

Phruriastis meliphaga, n. sp.

♂ ♀. 6–8 mm. Head dark grey, face whitish. Palpi dark fuscous, apex of second joint whitish, terminal joint ochreous-yellowish except base. Thorax grey, posterior edge whitish, shoulders and anterior edge dark grey. Forewings rather narrow, apex pointed, termen very obliquely rounded; pale grey; an elongate black spot on base of costa; small black spots on costa at $\frac{2}{5}$ and $\frac{3}{5}$, at tornus, and near apex: cilia grey-whitish. Hindwings and cilia grey.

FIJI, Labasa, September, October (*Veitch*); 3 ex. (type Brit. Mus.).

Catalectis ptilozona, n. sp.

♂ ♀. 8 mm. Head, thorax white. Labial palpi white, second joint dark fuscous except apex. Maxillary palpi blackish. Forewings white; an elongate dark brown spot on base of costa; a moderate somewhat oblique dark brown median fascia, slightly narrower towards costa, edged anteriorly by a ridge of raised scales, and posteriorly edged blackish and on lower half with a strong ridge-tuft grey anteriorly and blackish posteriorly; a small blackish dot on costa at $\frac{3}{4}$, and one in disc beneath it; a small metallic-grey spot at apex: cilia ochreous-whitish. Hindwings rather dark grey; cilia grey.

FIJI, Labasa, September, October (*Veitch*); 2 ex. (type Brit. Mus.).

Decadarchis subridens, n. sp.

♂. 10 mm. Head, palpi ochreous-whitish. Thorax whitish, anteriorly infuscated, a short expansible ochreous-whitish hair-pencil from beneath forewings. Forewings apex upturned; ochreous-white; markings dark brown; a mark on base of costa; an oblique spot from costa at $\frac{1}{4}$; an oblique mark from costa before middle uniting at a very acute angle with a very oblique fascia from $\frac{1}{3}$ of dorsum interrupted in disc, from angle of this an ochreous-yellow line runs just beneath costa to $\frac{5}{6}$, confluent with a very oblique striga from a dark fuscous mark on costa at $\frac{2}{3}$; an extremely oblique narrow slightly upcurved fascia running from $\frac{2}{3}$ of dorsum to apex, suffusedly interrupted in middle, an apical ochreous dot preceded by a black dot: cilia white, on costal origin a dark brown spot, above apex a fine dark fuscous hook. Hind-